

IMMUNIZATIONS / TESTS ONE OF THE FOLLOWING CRITERIA MUST BE MET: If documentation is not provided, the immunizations will be given (MMR is the vaccine of choice. If recipient is known to be immune to one or more of the components, monovalent or bivalent vaccines may be used)

Table: Evidence of Immunity to Vaccine Preventable Diseases and Tuberculosis

Disease	Evidence of Immunity/Screening	Immunization
Rubeola (Measles)	<p><u>Born before 1957:</u></p> <ol style="list-style-type: none"> 1. Documentation of one dose of live-measles vaccine <p style="text-align: center;"><i>or</i></p> <ol style="list-style-type: none"> 2. Documentation of a positive titer. <p><u>Born during or after 1957:</u></p> <ol style="list-style-type: none"> 1. Documentation of two doses of the vaccine after 1/1/1968 and on or after 1st birthday. <p style="text-align: center;"><i>or</i></p> <ol style="list-style-type: none"> 2. Documentation of a positive titer. 	<p>MMR will be given. 2nd dose given at least one month later.</p> <p>One dose of MMR</p>
Mumps	<p><u>Born before 1957:</u></p> <ol style="list-style-type: none"> 1. Documentation of one dose of live-mumps vaccine. <p><u>Born during or after 1957:</u></p> <ol style="list-style-type: none"> 1. Documentation of two doses of live mumps vaccine. The first dose of mumps containing vaccine administered on or after the 1st birthday. <p style="text-align: center;"><i>or</i></p> <ol style="list-style-type: none"> 2. Documentation of a positive titer. 	<p>One dose of live mumps virus vaccine.</p> <p>MMR will be given. 2nd dose given at least one month later.</p>
Rubella (German Measles)	<ol style="list-style-type: none"> 1. Documentation of one dose of the vaccine after 1/1/1970 and after 1st birthday. <p style="text-align: center;"><i>or</i></p> <ol style="list-style-type: none"> 2. Documentation of a positive titer 	<p>One dose of MMR</p>
Tuberculin Skin Test (TST)	<ol style="list-style-type: none"> 1. Documentation of TST <p style="text-align: center;"><i>or</i></p> <ol style="list-style-type: none"> 2. QuantiFERON[®]-TB Gold blood test* one year prior to screening date. <p style="text-align: center;"><i>or</i></p> <ol style="list-style-type: none"> 3. Documentation of a positive TST, with <i>either</i> <ol style="list-style-type: none"> a. Documentation of completion of INH treatment <p style="text-align: center;"><i>or</i></p> <ol style="list-style-type: none"> b. Documentation of a negative chest x-ray report. 	<p>TST repeated annually.</p>

Disease	Evidence of Immunity/Screening	Immunization
<p>Tetanus, Diphtheria and Pertusis (Tdap)</p> <p>Tetanus, Diphtheria</p>	<p>Documentation of vaccination.</p> <p>Documentation of vaccination.</p>	<p>A single dose of Tdap **</p> <p>A decennial booster with Td beginning 10 years after receipt of Tdap.</p>
<p>Varicella (chickenpox)</p>	<p>1. Documentation of 2 doses of vaccine.</p> <p><i>or</i></p> <p>2. Verbal history of the disease</p> <p><i>or</i></p> <p>3. Documentation of a positive titer.</p>	<p>For those with an uncertain history of disease a titer will be done. Varicella vaccine will be provided to susceptible health care workers. Two doses, four to eight weeks apart.</p>
<p>Influenza</p>	<p>Documentation of vaccination.</p>	<p>One dose annually.</p>
<p>Hepatitis B For healthcare workers who are identified as at risk for exposure to blood and body fluids (see Exposure Control Plan for Bloodborne Pathogens, Attachment I)</p>	<p>1. Documentation of hepatitis B vaccination series.</p> <p><i>or</i></p> <p>2. Documentation of a positive antibody titer.</p> <p><i>or</i></p> <p>3. History of disease (positive hepatitis B surface antigen).</p>	<p>Three dose series, two doses four weeks apart; 3rd dose five months after 2nd.</p>

* QuantiFERON[®]-TB Gold is not available in UEHC.

** Although Td booster doses are routinely recommended at an interval of 10 years, an interval as short as 2 years from the 1st dose of Td is recommended for HCW's who have direct patient contact. All other HCW's should receive a single dose of Tdap to replace the next scheduled Td.

- MMR is the vaccine of choice. If the recipient is known to be immune to one or more of the components, monovalent or bivalent vaccines may be used.
- Documentation is defined as a photocopy of the medical record or an immunization record signed by the health care provider or previous employer.